

t44_rpr_1
(TMb4DByEbx2UpZtsb4pnGXvfo8J7FcrkmB7)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_rpr_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0 X0) \wedge (v1_finset_1 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k2_rpr_1 X0 X1 (k3_subset_1 \\ & X0 X1) = k6_numbers)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k3_subset_1 \\ & X0 (k3_subset_1 X0 X1) = X1) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (m1_subset_1 \\ & (k3_subset_1 X0 X1) (k1_zfmisc_1 X0)) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0 X0) \wedge (v1_finset_1 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k2_rpr_1 X0 (k3_subset_1 X0 \\ & X1) X1 = k6_numbers)) \end{aligned}$$