

t44_sheffer2

(TMQneC11BvwhoJfyNqX3gLw6qEqADqCZcUg)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v1_sheffer2 : \iota \Rightarrow o$ be given. Let $l1_sheffer1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k5_sheffer1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & \quad X0))) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 X1 (k5_sheffer1 \\ & \quad X0 X2 X1) = k5_sheffer1 X0 X1 (k5_sheffer1 X0 X2 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & \quad X0))) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 X1 X2 = k5_sheffer1 \\ & \quad X0 X2 X1))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & \quad X0))) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 X1 (k5_sheffer1 \\ & \quad X0 X2 X2) = k5_sheffer1 X0 X1 (k5_sheffer1 X0 X1 X2)))) \end{aligned}$$