

t44\_xcmlx\_1

(TMc2MwMUUjHi5bsFH8RBi7dpvkEQMSjvQo3)

October 27, 2020

Let  $v1\_xcmlx\_0 : \iota \Rightarrow o$  be given. Let  $k3\_xcmlx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmlx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_xcmlx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xcmlx\_0 X0) \Rightarrow (\forall X1.(v1\_xcmlx\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xcmlx\_0 X2) \Rightarrow (\forall X3.(v1\_xcmlx\_0 X3) \Rightarrow (k3\_xcmlx\_0 \\ & (k6\_xcmlx\_0 (k2\_xcmlx\_0 X0 X1) X2) X3 = k6\_xcmlx\_0 (k2\_xcmlx\_0 \\ & (k3\_xcmlx\_0 X0 X3) (k3\_xcmlx\_0 X1 X3)) (k3\_xcmlx\_0 X2 X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xcmlx\_0 X0) \Rightarrow (\forall X1.(v1\_xcmlx\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xcmlx\_0 X2) \Rightarrow (k6\_xcmlx\_0 (k2\_xcmlx\_0 X0 X1) X2 = k2\_xcmlx\_0 \\ & (k6\_xcmlx\_0 X0 X2) X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1\_xcmlx\_0 X0) \wedge (v1\_xcmlx\_0 X1)) \Rightarrow ( \\ & v1\_xcmlx\_0 (k3\_xcmlx\_0 X0 X1)) \end{aligned} \quad (3)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(v1\_xcmlx\_0 X0) \Rightarrow (\forall X1.(v1\_xcmlx\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xcmlx\_0 X2) \Rightarrow (\forall X3.(v1\_xcmlx\_0 X3) \Rightarrow (k3\_xcmlx\_0 \\ & (k2\_xcmlx\_0 (k6\_xcmlx\_0 X0 X1) X2) X3 = k2\_xcmlx\_0 (k6\_xcmlx\_0 \\ & (k3\_xcmlx\_0 X0 X3) (k3\_xcmlx\_0 X1 X3)) (k3\_xcmlx\_0 X2 X3)))))) \end{aligned}$$