

t44_xreal_1 (TMdEy-
cdQ2VBHweXkzV5t54U3dP5RbNPoRhU)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge (r1_xxreal_0 (k2_xcmplx_0 X1 X0) X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 (k6_xcmplx_0 X1 X2)) \Rightarrow (r1_xxreal_0 (k2_xcmplx_0 X0 X2) X1)))) \quad (2)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge (r1_xxreal_0 X1 (k6_xcmplx_0 X1 X0))))$$