

t45_circcomb
(TMPt4jdKSs6TJBjFRE3GMzQe8vFmZM3hSoH)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_msualg_1 : \iota \Rightarrow o$ be given. Let $v1_circcomb : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u4_struct_0 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k6_partfun1 : \iota \Rightarrow \iota$ be given. Let $k4_relat_1 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $u2_msualg_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski X1 X2)) \Rightarrow (r1_tarski X0 X2) \quad (1)$$

Assume the following.

$$\forall X0. k6_partfun1 X0 = k4_relat_1 X0 \quad (2)$$

Assume the following.

$$\forall X0. k10_xtuple_0 (k4_relat_1 X0) = X0 \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0. (l1_msualg_1 X0) \Rightarrow & ((v1_funct_1 (u2_msualg_1 X0)) \wedge \\ & ((v1_funct_2 (u2_msualg_1 X0) (u4_struct_0 X0) (u1_struct_0 X0)) \wedge \\ & (m1_subset_1 (u2_msualg_1 X0) (k1_zfmisc_1 (k2_zfmisc_1 (u4_struct_0 \\ & X0) (u1_struct_0 X0)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0. v1_relat_1 (k4_relat_1 X0) \quad (5)$$

Assume the following.

$$\forall X0. (l1_msualg_1 X0) \Rightarrow ((v1_circcomb X0) \Leftrightarrow (u2_msualg_1 X0 = k6_partfun1 (u4_struct_0 X0))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X1)\Rightarrow((v5_relat_1 X1 X0)\Leftrightarrow(r1_tarski (k10_xtuple_0 X1) X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(X0 = X1)\Leftrightarrow((r1_tarski X0 X1)\wedge(r1_tarski X1 X0)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (9)$$

Theorem 1

$$\forall X0.((\neg v2_struct_0 X0)\wedge(l1_msualg_1 X0))\Rightarrow((v1_circcomb X0)\Rightarrow(r1_tarski (u4_struct_0 X0) (u1_struct_0 X0)))$$