

t45_funcop_1 (TMQJzuTM- CNh1txNCnAMAvwEAnLBEPcSRcLp)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (X1 \in X0) \Rightarrow (k1_funct_1 (k2_funcop_1 X0 X2) X1 = X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow (((k9_xtuple_0 X2 = X0) \wedge (\forall X3. (X3 \in X0) \Rightarrow (k1_funct_1 X2 X3 \in X1))) \Rightarrow ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k2_funcop_1 X0 X1)) \wedge (v1_funct_1 (k2_funcop_1 X0 X1)) \quad (4)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. (X2 \in X1) \Rightarrow ((v1_funct_1 (k2_funcop_1 X0 X2)) \wedge ((v1_funct_2 (k2_funcop_1 X0 X2) X0 X1) \wedge (m1_subset_1 (k2_funcop_1 X0 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))))$$