

t45_sheffer2 (TMHd-
sCb3FXkoMVEtGH8JQQegi1W5DqA7w5P)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v1_sheffer2 : \iota \Rightarrow o$ be given. Let $l1_sheffer1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k5_sheffer1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 (k5_sheffer1 \\ & X0 X1 X2) X2 = k5_sheffer1 X0 X2 (k5_sheffer1 X0 X1 X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 \\ & (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 (k5_sheffer1 X0 (k5_sheffer1 \\ & X0 X1 X2) X2) (k5_sheffer1 X0 X2 (k5_sheffer1 X0 X3 X1)) = k5_sheffer1 \\ & X0 (k5_sheffer1 X0 X2 (k5_sheffer1 X0 X3 X1)) (k5_sheffer1 X0 X2 (\\ & k5_sheffer1 X0 X3 X1)))))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 \\ & (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 (k5_sheffer1 X0 X1 (k5_sheffer1 \\ & X0 X2 X2)) (k5_sheffer1 X0 X1 (k5_sheffer1 X0 X3 X2)) = k5_sheffer1 \\ & X0 (k5_sheffer1 X0 X1 (k5_sheffer1 X0 X3 X2)) (k5_sheffer1 X0 X1 (\\ & k5_sheffer1 X0 X3 X2)))))) \end{aligned}$$