

t46_bvfunc14

(TMMg7ZcpEJmDvf5PCGQ1k4zA8mCPLnFVU2Q)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_bvfunc_2 : \iota \Rightarrow \iota$ be given. Let $m1_eqrel_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_bvfunc_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_partit1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.k2_enumset1\ X0\ X1 \\ X2\ X3 = k2_xboole_0\ (k2_tarski\ X0\ X1)\ (k2_tarski\ X2\ X3) \quad (1)$$

Assume the following.

$$\forall X0.(\neg v1_xboole_0\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1 \\ (k1_bvfunc_2\ X0))) \Rightarrow (\forall X2.(m1_eqrel_1\ X2\ X0) \Rightarrow (\forall X3. \\ (m1_eqrel_1\ X3\ X0) \Rightarrow (\forall X4.(m1_eqrel_1\ X4\ X0) \Rightarrow (\forall X5. \\ (m1_eqrel_1\ X5\ X0) \Rightarrow (\forall X6.(m1_eqrel_1\ X6\ X0) \Rightarrow (\forall X7. \\ (m1_eqrel_1\ X7\ X0) \Rightarrow (\forall X8.(m1_eqrel_1\ X8\ X0) \Rightarrow ((X1 = k5_enumset1 \\ X2\ X3\ X4\ X5\ X6\ X7\ X8) \Rightarrow ((X2 = X3) \vee ((X2 = X4) \vee ((X2 = X5) \vee ((X2 = X6) \vee ((X2 = \\ X7) \vee ((X2 = X8) \vee ((X3 = X4) \vee ((X3 = X5) \vee ((X3 = X6) \vee ((X3 = X7) \vee ((X3 = X8) \vee \\ ((X4 = X5) \vee ((X4 = X6) \vee ((X4 = X7) \vee ((X4 = X8) \vee ((X5 = X6) \vee ((X5 = X7) \vee \\ (X5 = X8) \vee ((X6 = X7) \vee ((X6 = X8) \vee ((X7 = X8) \vee (k5_bvfunc_2\ X0\ X5\ X1 = k2_partit1 \\ X0\ (k2_partit1\ X0\ (k2_partit1\ X0\ (k2_partit1\ X0\ (k2_partit1\ X0\ X2 \\ X3)\ X4)\ X6)\ X7)\ X8)))))))))))))))))))))))))))))))))))))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.\forall X5. \\ \forall X6.k5_enumset1\ X0\ X1\ X2\ X3\ X4\ X5\ X6 = k2_xboole_0\ (k1_enumset1 \\ X0\ X1\ X2)\ (k2_enumset1\ X3\ X4\ X5\ X6) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.k2_tarski\ X0\ X1 = k2_tarski\ X1\ X0 \quad (4)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ & \quad (k1_bfunc_2 X0))) \Rightarrow (\forall X2.(m1_eqrel_1 X2 X0) \Rightarrow (\forall X3. \\ & \quad (m1_eqrel_1 X3 X0) \Rightarrow (\forall X4.(m1_eqrel_1 X4 X0) \Rightarrow (\forall X5. \\ & \quad (m1_eqrel_1 X5 X0) \Rightarrow (\forall X6.(m1_eqrel_1 X6 X0) \Rightarrow (\forall X7. \\ & \quad (m1_eqrel_1 X7 X0) \Rightarrow (\forall X8.(m1_eqrel_1 X8 X0) \Rightarrow ((X1 = k5_enumset1 \\ & X2 X3 X4 X5 X6 X7 X8) \Rightarrow ((X2 = X3) \vee ((X2 = X4) \vee ((X2 = X5) \vee ((X2 = X6) \vee ((X2 = \\ & X7) \vee ((X2 = X8) \vee ((X3 = X4) \vee ((X3 = X5) \vee ((X3 = X6) \vee ((X3 = X7) \vee ((X3 = X8) \vee \\ & ((X4 = X5) \vee ((X4 = X6) \vee ((X4 = X7) \vee ((X4 = X8) \vee ((X5 = X6) \vee ((X5 = X7) \vee (\\ & (X5 = X8) \vee ((X6 = X7) \vee ((X6 = X8) \vee ((X7 = X8) \vee (k5_bfunc_2 X0 X6 X1 = k2_partit1 \\ & X0 (k2_partit1 X0 (k2_partit1 X0 (k2_partit1 X0 (k2_partit1 X0 X2 \\ & X3) X4) X5) X7) X8)) \end{aligned}$$