

t46_coh_sp (TMQEF-
fqJ3fo5B8WYmh7BHoPqk1rqbsNmZua)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k16_coh_sp : \iota \Rightarrow \iota$ be given. Let $k2_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k23_coh_sp : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_partfun1 : \iota \Rightarrow \iota$ be given. Let $k17_coh_sp : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k21_coh_sp : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k22_coh_sp : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k20_coh_sp : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. k2_xtuple_0 (k4_tarski X0 X1) = X1 \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k1_xtuple_0 (k4_tarski X0 X1) = X0 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k16_coh_sp X0)) \Rightarrow (m1_subset_1 (k23_coh_sp X0 X1) (k20_coh_sp X0)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k16_coh_sp X0)) \Rightarrow (k23_coh_sp X0 X1 = k4_tarski (k4_tarski X1 X1) (k6_partfun1 (k17_coh_sp X0 X1))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k20_coh_sp X0)) \Rightarrow (k22_coh_sp X0 X1 = k2_xtuple_0 (k1_xtuple_0 X1)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k20_coh_sp X0)) \Rightarrow (k21_coh_sp X0 X1 = k1_xtuple_0 (k1_xtuple_0 X1)) \quad (6)$$

Theorem 1

$$\forall X0. \forall X1. (m1_subset_1 X1 (k16_coh_sp X0)) \Rightarrow ((k2_xtuple_0 (k23_coh_sp X0 X1) = k6_partfun1 (k17_coh_sp X0 X1)) \wedge ((k21_coh_sp X0 (k23_coh_sp X0 X1) = X1) \wedge (k22_coh_sp X0 (k23_coh_sp X0 X1) = X1)))$$