

t46\_eqrel\_1  
(TMG1eLvWu3xqruHa5aNPKh1me9QnKxp7KFB)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k8\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (r1\_tarski (k7\_relat\_1 X1 (k8\_relat\_1 X1 X0)) X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1\_tarski X0 X1) \wedge (r1\_tarski X1 X2)) \Rightarrow (r1\_tarski X0 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (v1\_relat\_1 X2) \Rightarrow ((r1\_tarski X0 X1) \Rightarrow (r1\_tarski (k7\_relat\_1 X2 X0) (k7\_relat\_1 X2 X1))) \quad (3)$$

**Theorem 1**

$$\forall X0. \forall X1. \forall X2. ((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow ((r1\_tarski X0 (k8\_relat\_1 X2 X1)) \Rightarrow (r1\_tarski (k7\_relat\_1 X2 X0) X1))$$