

t46\_setfam\_1  
(TMPp8BiFNLckTr3dRC5tWwSkUYn4ALffKg4)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_tarSKI : \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 X0) \Rightarrow ((X0 \neq k1\_xboole\_0) \Rightarrow (m1\_subset\_1 (k1\_tarSKI X1) (k1\_zfmisc\_1 X0))) \quad (1)$$

Assume the following.

$$\forall X0. m1\_subset\_1 k1\_xboole\_0 (k1\_zfmisc\_1 X0) \quad (2)$$

Assume the following.

$$v1\_xboole\_0 k1\_xboole\_0 \quad (3)$$

Assume the following.

$$\forall X0. \neg v1\_xboole\_0 (k1\_zfmisc\_1 X0) \quad (4)$$

**Theorem 1**

$$\forall X0. m1\_subset\_1 (k1\_tarSKI k1\_xboole\_0) (k1\_zfmisc\_1 (k1\_zfmisc\_1 X0))$$