

t47_facirc_1 (TMZYCfsekmUKdgHauHinjJKDur- GAcaWhEcg)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_msafree2 : \iota \Rightarrow \iota$ be given. Let $k5_circcomb : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_msafree2 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ X1))) \Rightarrow ((k2_msafree2 (k5_circcomb X0 X1) = k10_xtuple_0 X1) \wedge (k3_msafree2 \\ (k5_circcomb X0 X1) = k1_tarski (k4_tarski X1 X0))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \tag{2}$$

Theorem 1

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (\forall X1. k4_tarski X0 X1 \in k3_msafree2 (k5_circcomb X1 X0)) \end{aligned}$$