

t47_group_2
(TMdCC9xfUBR5CYTqCr9yM6jVBezqxSbXi9j)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_group_1 : \iota \Rightarrow o$ be given. Let $v3_group_1 : \iota \Rightarrow o$ be given. Let $l3_algstr_0 : \iota \Rightarrow o$ be given. Let $m1_group_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_struct_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_group_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.(m1_group_2 X1 X0) \Rightarrow (r1_struct_0 X1 (k1_group_1 X0))) \quad (1)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.(m1_group_2 X1 X0) \Rightarrow (k1_group_1 X1 = k1_group_1 X0)) \quad (2)$$

Theorem 1

$$\forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.(m1_group_2 X1 X0) \Rightarrow (\forall X2.(m1_group_2 X2 X0) \Rightarrow (r1_struct_0 X2 (k1_group_1 X1))))$$