

t47_hilbert1 (TMdzN-
brKyJ9wcv96BNQzwTHx98hAvbRkVJ8)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_hilbert1 : \iota$ be given. Let $k3_hilbert1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_hilbert1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_hilbert1 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k1_hilbert1) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k1_hilbert1) \Rightarrow (\forall X2.(m1_subset_1 X2 k1_hilbert1) \Rightarrow (k3_hilbert1 \\ & (k3_hilbert1 X0 X1) (k3_hilbert1 (k4_hilbert1 X2 X0) (k4_hilbert1 \\ & X2 X1)) \in k6_hilbert1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k1_hilbert1) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k1_hilbert1) \Rightarrow (k3_hilbert1 (k4_hilbert1 X0 X1) (k4_hilbert1 \\ & X1 X0) \in k6_hilbert1)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k1_hilbert1) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k1_hilbert1) \Rightarrow (\forall X2.(m1_subset_1 X2 k1_hilbert1) \Rightarrow ((\\ & (k3_hilbert1 X0 (k3_hilbert1 X1 X2) \in k6_hilbert1) \wedge (X1 \in k6_hilbert1)) \Rightarrow \\ & (k3_hilbert1 X0 X2 \in k6_hilbert1)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k1_hilbert1) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k1_hilbert1) \Rightarrow (\forall X2.(m1_subset_1 X2 k1_hilbert1) \Rightarrow ((\\ & k3_hilbert1 X0 (k3_hilbert1 X1 X2) \in k6_hilbert1) \Rightarrow (k3_hilbert1 \\ & X1 (k3_hilbert1 X0 X2) \in k6_hilbert1)))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1_subset_1 X0 k1_hilbert1) \wedge (m1_subset_1 \\ & X1 k1_hilbert1)) \Rightarrow (m1_subset_1 (k4_hilbert1 X0 X1) k1_hilbert1) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1_subset_1 X0 k1_hilbert1) \wedge (m1_subset_1 \\ & X1 k1_hilbert1)) \Rightarrow (m1_subset_1 (k3_hilbert1 X0 X1) k1_hilbert1) \end{aligned} \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k1_hilbert1) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k1_hilbert1) \Rightarrow (\forall X2.(m1_subset_1 X2 k1_hilbert1) \Rightarrow (k3_hilbert1 \\ & (k4_hilbert1 X0 (k4_hilbert1 X1 X2)) (k4_hilbert1 X0 (k4_hilbert1 \\ & X2 X1)) \in k6_hilbert1))) \end{aligned}$$