

t47_mesfunc6

(TMNbqQ4q4es9yvuaF3x5mekqMJ14UiHkwyD)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_mesfunc6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k19_rfunct_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_mesfunc2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_mesfunc5 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_rfunct_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_mesfunc2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_numbers : \iota$ be given. Let $r1_mesfunc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ (k1_mesfunc2 X0 (k1_mesfunc5 X0 X1) = k18_rfunct_3 X0 X1) \wedge (k2_mesfunc2 \\ X0 (k1_mesfunc5 X0 X1) = k19_rfunct_3 X0 X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \Rightarrow (\\ \forall X2. ((\neg v1_xboole_0 X2) \wedge ((v1_prob_1 X2 X0) \wedge ((v4_prob_1 \\ X2 X0) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X3. \\ (m1_subset_1 X3 X2) \Rightarrow (((r1_mesfunc1 X0 X2 X1 X3) \wedge (r1_tarski X3 (\\ k1_relset_1 X0 X1))) \Rightarrow (r1_mesfunc1 X0 X2 (k2_mesfunc2 X0 X1) X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \neg v1_xboole_0 (k1_zfmisc_1 X0) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge \\ & m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow \\ & ((v1_funct_1 (k1_mesfunc5 X0 X1)) \wedge (m1_subset_1 (k1_mesfunc5 \\ & X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge \\ & m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow \\ & ((v1_funct_1 (k19_rfunct_3 X0 X1)) \wedge (m1_subset_1 (k19_rfunct_3 \\ & X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge \\ & m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ & k1_mesfunc5 X0 X1 = X1) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\forall X3. (\\ & m2_subset_1 X3 (k1_zfmisc_1 X0) X1) \Rightarrow ((r1_mesfunc6 X0 X1 X2 X3) \Leftrightarrow \\ & (r1_mesfunc1 X0 X1 (k1_mesfunc5 X0 X2) X3)))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\forall X3. (\\ & m2_subset_1 X3 (k1_zfmisc_1 X0) X1) \Rightarrow (((r1_mesfunc6 X0 X1 X2 X3) \wedge \\ & (r1_tarSKI X3 (k1_relset_1 X0 X2))) \Rightarrow (r1_mesfunc6 X0 X1 (k19_rfunct_3 \\ & X0 X2) X3)))) \end{aligned}$$