

t47_rlvect_1
(TMZr8ZimqzXzYpiJj4kEeX6KbCc6RJPQski)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_finseq_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k4_struct_0 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $l2_algstr_0 : \iota \Rightarrow o$ be given. Let $l2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_algstr_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.((\neg v2_struct_0 X1) \wedge (\\ v13_algstr_0 X1) \wedge ((v2_rlvect_1 X1) \wedge ((v3_rlvect_1 X1) \wedge ((v4_rlvect_1 \\ X1) \wedge ((v5_rlvect_1 X1) \wedge ((v6_rlvect_1 X1) \wedge ((v7_rlvect_1 X1) \wedge \\ ((v8_rlvect_1 X1) \wedge (l1_rlvect_1 X1)))))))))) \Rightarrow (\forall X2.(m1_subset_1 \\ X2 (u1_struct_0 X1)) \Rightarrow (((X0 = k6_numbers) \vee (X2 = k4_struct_0 X1)) \Rightarrow \\ (k1_rlvect_1 X1 X2 X0 = k4_struct_0 X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l2_algstr_0 X0)) \Rightarrow (k4_rlvect_1 X0 (k6_finseq_1 (u1_struct_0 X0)) = k4_struct_0 X0) \tag{2}$$

Assume the following.

$$\forall X0.(l2_algstr_0 X0) \Rightarrow ((l2_struct_0 X0) \wedge (l1_algstr_0 X0)) \tag{3}$$

Assume the following.

$$\forall X0.(l1_rlvect_1 X0) \Rightarrow (l2_algstr_0 X0) \tag{4}$$

Assume the following.

$$\forall X0.(l2_struct_0 X0) \Rightarrow (m1_subset_1 (k4_struct_0 X0) (u1_struct_0 X0)) \tag{5}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\ & X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\ & X0)))))))))) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow (k1_rlvect_1 \\ & X0 (k4_rlvect_1 X0 (k6_finseq_1 (u1_struct_0 X0))) X1 = k4_struct_0 \\ & X0)) \end{aligned}$$