

t48_card_3

(TMQwAYfJmH8oHCAqDc93idtGC7x8Z4HHbBt)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k4_card_3 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(X1 = \\ k4_card_3 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.((v1_relat_1 \\ X3) \wedge (v1_funct_1 X3)) \wedge ((X2 = X3) \wedge ((k9_xtuple_0 X3 = k9_xtuple_0 \\ X0) \wedge (\forall X4.(X4 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 X3 X4 \in k1_funct_1 \\ X0 X4))))))) \end{aligned} \tag{1}$$

Theorem 1

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((X0 \in k4_card_3 X1) \Rightarrow ((v1_relat_1 X0) \wedge (v1_funct_1 X0)))$$