

t48_xboolean (TMPDNU- JisK6qSRgY45fXtZ8UkVaNYD1L4GB)

October 27, 2020

Let $v1_xboolean : \iota \Rightarrow o$ be given. Let $k4_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboolean : \iota \Rightarrow \iota$ be given. Let $k7_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k9_xboolean X0 (k5_xboolean X0 X1) = k9_xboolean X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k9_xboolean X0 (k5_xboolean X0 X1) = k4_xboolean (k3_xboolean X0) (k3_xboolean X1))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k4_xboolean (k3_xboolean X0) (k7_xboolean X0 X1) = k4_xboolean (k3_xboolean X0) (k3_xboolean X1))) \quad (3)$$

Theorem 1

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k4_xboolean (k3_xboolean X0) (k7_xboolean X0 X1) = k9_xboolean X0 X1))$$