

t48\_xxreal\_0  
(TMbS7iVSLriMLVmJv1Hwu3zn1eeDNFjUBRk)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k2\_xxreal\_0 : \iota$  be given. Let  $k1\_xxreal\_0 : \iota$  be given. Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (r1\_xxreal\_0 k2\_xxreal\_0 X0) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (r1\_xxreal\_0 X0 k1\_xxreal\_0) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\neg(\neg X0 \in k1\_numbers) \wedge ((X0 \neq k1\_xxreal\_0) \wedge (X0 \neq k2\_xxreal\_0))) \quad (3)$$

Assume the following.

$$k1\_xxreal\_0 = k1\_numbers \quad (4)$$

**Theorem 1**

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2.(v1\_xxreal\_0 X2) \Rightarrow (\neg(\neg r1\_xxreal\_0 X1 X0) \wedge ((\neg r1\_xxreal\_0 X2 X1) \wedge (\neg X1 \in k1\_numbers))))))$$