

t49_bvfunc_1

(TMarsnsAFZ2og6FVf1EP5wBbqFaj35ad6tE)

October 27, 2020

Let $v1_xboolean : \iota \Rightarrow o$ be given. Let $k8_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboolean : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (\forall X2. \\ & (v1_xboolean X2) \Rightarrow (k8_xboolean X0 (k4_xboolean X1 X2) = k3_xboolean \\ & (k4_xboolean (k4_xboolean X0 X1) X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (v1_xboolean (k4_xboolean X0 X1)) \tag{2}$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k8_xboolean X0 X1 = k3_xboolean (k4_xboolean X0 X1))) \tag{3}$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (\forall X2. \\ & (v1_xboolean X2) \Rightarrow (k8_xboolean X0 (k4_xboolean X1 X2) = k8_xboolean \\ & (k4_xboolean X0 X1) X2)))) \end{aligned}$$