

t49_finseq_6

(TMRh4p4JytM8Y3PL8U3Z5d6JLoTSWrJoRyY)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_finseq_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow \\ & (\forall X2. (m2_finseq_1 X2 X0) \Rightarrow ((X1 \in k10_xtuple_0 X2) \Rightarrow (k1_finseq_5 \\ & X0 X2 X1 = k7_finseq_1 (k5_finseq_4 X2 X1) (k12_finseq_1 X0 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \neq X1) \Rightarrow (k5_finseq_4 (k10_finseq_1 X0 X1) X1 = k9_finseq_1 X0) \quad (2)$$

Assume the following.

$$\forall X0. k9_finseq_1 X0 = k5_finseq_1 X0 \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((\neg v1_xboole_0 X0) \wedge ((m1_subset_1 \\ & X1 X0) \wedge (m1_subset_1 X2 X0))) \Rightarrow (k2_finseq_4 X0 X1 X2 = k10_finseq_1 \\ & X1 X2) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k12_finseq_1 X0 X1 = k5_finseq_1 X1) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. k10_xtuple_0 (k10_finseq_1 X0 X1) = k2_tarski X0 X1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge((m1_subset_1 X1 X0)\wedge(m1_subset_1 X2 X0)))\Rightarrow(m2_finseq_1 (k2_finseq_4 X0 X1 X2) X0) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k10_finseq_1 X0 X1 = k7_finseq_1 (k9_finseq_1 X0) (k9_finseq_1 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k2_tarSKI X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 = X0)\vee(X3 = X1))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.k2_tarSKI X0 X1 = k2_tarSKI X1 X0 \quad (10)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(\forall X2.(m1_subset_1 X2 X0)\Rightarrow((X1\neq X2)\Rightarrow(k1_finseq_5 X0 (k2_finseq_4 X0 X1 X2) X2 = k2_finseq_4 X0 X1 X2))))$$