

t49_flang_2
(TMKfh4aVobTrwSbja8ib7LY4PC8vxFKc8gn)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $k1_flang_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $np_2 : \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_catalan2 : \iota \Rightarrow \iota$ be given. Let $k6_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & X0))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (k1_flang_2 X0 X1 X2 (k2_xcmplx_0 \\ & X2 np_1) = k4_subset_1 (k3_catalan2 X0) (k7_flang_1 X0 X1 X2) (k7_flang_1 \\ & X0 X1 (k2_xcmplx_0 X2 np_1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k3_catalan2 X0))) \Rightarrow (k7_flang_1 X0 X1 np_2 = k6_flang_1 X0 X1 X1) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k3_catalan2 X0))) \Rightarrow (k7_flang_1 X0 X1 np_1 = X1) \tag{3}$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \tag{4}$$

Assume the following.

$$k2_xcmplx_0 np_1 np_1 = np_2 \tag{5}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{6}$$

Assume the following.

$$\forall X0.k3_catalan2\ X0 = k8_afinsq_1\ X0 \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1) \Rightarrow (v7_ordinal1\ X0) \quad (8)$$

Theorem 1

$$\forall X0.\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ (k8_afinsq_1\ X0))) \Rightarrow (k1_flang_2\ X0\ X1\ np_1\ np_2 = k4_subset_1\ (k3_catalan2\ X0)\ X1\ (k6_flang_1\ X0\ X1\ X1))$$