

t49_funct_1 (TM-
SPofAGx7mb5CoQvQgRx3KD2viKSuMZLxj)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((X0 \in k9_xtuple_0 (k5_relat_1 X2 X1)) \Leftrightarrow ((X0 \in X1) \wedge (X0 \in k9_xtuple_0 X2))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 (k5_relat_1 X2 X1)) \Rightarrow (k1_funct_1 (k5_relat_1 X2 X1) X0 = k1_funct_1 X2 X0)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v1_relat_1 (k5_relat_1 X0 X1)) \wedge (v1_funct_1 (k5_relat_1 X0 X1))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X0) \Rightarrow (v1_relat_1 (k5_relat_1 X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. ((X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (k4_tarski X1 X2 \in X0))) \wedge ((\neg X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in X1) \Rightarrow (k1_funct_1 (k5_relat_1 X2 X1) X0 = k1_funct_1 X2 X0))$$