

t49_partfun2

(TMaoNyDe96YBhRJFcUiWyuPkRNeKkEweSjL)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((k9_xtuple_0 \\ & X1 = k1_tarski X0) \Rightarrow (X1 = k1_tarski (k4_tarski X0 (k1_funct_1 X1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (\\ & k1_relset_1 X0 X1 = k9_xtuple_0 X1) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v5_relat_1 X1 X0) \wedge (\\ & v1_funct_1 X1))) \Rightarrow (\forall X2. (X2 \in k9_xtuple_0 X1) \Rightarrow (k7_partfun1 \\ & X0 X1 X2 = k1_funct_1 X1 X2)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow \\ & (X2 = X0)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \end{aligned} \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow \\ & (\forall X2.(m1_subset_1 X2 X0) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge \\ & (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow ((k1_relset_1 \\ & X0 X3 = k1_tarski X2) \Rightarrow (X3 = k1_tarski (k4_tarski X2 (k7_partfun1 \\ & X1 X3 X2))))))) \end{aligned}$$