

t49_power
(TMVPprYSD3Zfqwe1Jh2Wh7ohotGa59zfdgS)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_power : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\neg \\ r1_xxreal_0 X0 (k1_real_1 np_1)) \Rightarrow (r1_xxreal_0 (k7_real_1 np_1 \\ (k3_xcmplx_0 X1 X0)) (k2_newton (k7_real_1 np_1 X0) X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v7_ordinal1 X1)) \Rightarrow (k2_newton X0 X1 = k1_newton X0 X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (k3_power X0 X1 = k1_newton X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k7_real_1 X0 X1) k1_numbers) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 k5_numbers) \Rightarrow \\ & ((\neg r1_xxreal_0 X0 (k1_real_1 np_1)) \Rightarrow (r1_xxreal_0 (k7_real_1 \\ & np_1 (k3_xcmplx_0 X1 X0)) (k3_power (k7_real_1 np_1 X0) X1)))) \end{aligned}$$