

t49_sheffer2 (TMauTMjQdMnnSeCuouovfqCN- MXeArvrM548)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v1_sheffer2 : \iota \Rightarrow o$ be given. Let $l1_sheffer1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k5_sheffer1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 \\ & (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 X1 (k5_sheffer1 X0 (k5_sheffer1 \\ & X0 X2 X2) (k5_sheffer1 X0 X3 (k5_sheffer1 X0 X1 (k5_sheffer1 X0 X1 \\ & X2)))) = k5_sheffer1 X0 X1 (k5_sheffer1 X0 (k5_sheffer1 X0 X2 X2) \\ & (k5_sheffer1 X0 X2 X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 (k5_sheffer1 \\ & X0 X1 X2) (k5_sheffer1 X0 X1 X1) = X1))) \end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 \\ & (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 X1 (k5_sheffer1 X0 (k5_sheffer1 \\ & X0 X2 X2) (k5_sheffer1 X0 X3 (k5_sheffer1 X0 X1 (k5_sheffer1 X0 X1 \\ & X2)))) = k5_sheffer1 X0 X1 X2)))) \end{aligned}$$