

# t49\_tops\_2 (TMUjoXcGCboyCT- MXLVK4KeYHTUJ43Cc4tki)

October 27, 2020

Let  $l1\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v2\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_tops\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k2\_funct\_1 : \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v3\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v2\_funct\_2 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow ((v2\_funct\_1 X0) \Rightarrow \\ ((k10\_xtuple\_0 X0 = k9\_xtuple\_0 (k2\_funct\_1 X0)) \wedge (k9\_xtuple\_0 \\ X0 = k10\_xtuple\_0 (k2\_funct\_1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v5\_relat\_1 X1 X0)) \Rightarrow (k2\_relset\_1 X0 X1 = k10\_xtuple\_0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (3)$$

Assume the following.

$$\forall X0. ((\neg v2\_struct\_0 X0) \wedge (l1\_struct\_0 X0)) \Rightarrow (\neg v1\_xboole\_0 (u1\_struct\_0 X0)) \quad (4)$$

Assume the following.

$$v1\_xboole\_0 k1\_xboole\_0 \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v1\_funct\_1 X2)\wedge((v1\_funct\_2 \\ & X2 X0 X1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))\Rightarrow \\ & ((v1\_funct\_1 (k2\_tops\_2 X0 X1 X2))\wedge((v1\_funct\_2 (k2\_tops\_2 X0 \\ & X1 X2) X1 X0)\wedge(m1\_subset\_1 (k2\_tops\_2 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X1 X0)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v1\_funct\_1 X2)\wedge((v1\_funct\_2 \\ & X2 X0 X1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))\Rightarrow \\ & ((v3\_funct\_2 X2 X0 X1)\Rightarrow(k2\_tops\_2 X0 X1 X2 = k2\_funct\_1 X2)) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0.(l1\_struct\_0 X0)\Rightarrow(k2\_struct\_0 X0 = u1\_struct\_0 X0) \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v5\_relat\_1 X1 X0))\Rightarrow( \\ & (v2\_funct\_2 X1 X0)\Leftrightarrow(k2\_relset\_1 X0 X1 = X0)) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))\Rightarrow(((X1\neq k1\_xboole\_0)\Rightarrow((v1\_funct\_2 X2 X0 \\ & X1)\Leftrightarrow(X0 = k1\_relset\_1 X0 X2)))\wedge((X1 = k1\_xboole\_0)\Rightarrow((v1\_funct\_2 \\ & X2 X0 X1)\Leftrightarrow(X2 = k1\_xboole\_0)))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))\Rightarrow(((v1\_funct\_1 X2)\wedge((v2\_funct\_1 X2)\wedge(v2\_funct\_2 \\ & X2 X1)))\Rightarrow((v1\_funct\_1 X2)\wedge(v3\_funct\_2 X2 X0 X1))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \end{aligned} \quad (13)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(l1\_struct\_0 X0) \Rightarrow (\forall X1.((\neg v2\_struct\_0 X1) \wedge \\ & (l1\_struct\_0 X1)) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\ & X2 (u1\_struct\_0 X0) (u1\_struct\_0 X1)) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 (u1\_struct\_0 X0) (u1\_struct\_0 X1)))))) \Rightarrow (((k2\_relset\_1 \\ & (u1\_struct\_0 X1) X2 = k2\_struct\_0 X1) \wedge (v2\_funct\_1 X2)) \Rightarrow ((k1\_relset\_1 \\ & (u1\_struct\_0 X1) (k2\_tops\_2 (u1\_struct\_0 X0) (u1\_struct\_0 X1) \\ & X2) = k2\_struct\_0 X1) \wedge (k2\_relset\_1 (u1\_struct\_0 X0) (k2\_tops\_2 \\ & (u1\_struct\_0 X0) (u1\_struct\_0 X1) X2) = k2\_struct\_0 X0)))))) \end{aligned}$$