

t49_xxreal_3 (TMNzMiU-
uyuHXQS8vtX9WyfMBPDpnWWsCMsU)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k1_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k3_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k5_numbers : \iota$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (k1_xxreal_3 X0 k6_numbers = X0) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ (v1_xreal_0 X2) \Rightarrow (\neg(\neg r1_xxreal_0 X1 X0) \wedge ((\neg r1_xxreal_0 X2 X1) \wedge \\ (\neg X1 \in k1_numbers)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg \\ (\neg r1_xxreal_0 X0 k6_numbers) \wedge ((\neg r1_xxreal_0 X1 X0) \wedge (r1_xreal_0 \\ (k3_xxreal_3 X1 X0) k6_numbers)))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg \\ (\neg r1_xxreal_0 X1 X0) \wedge (\forall X2.(v1_xreal_0 X2) \Rightarrow (\neg(\neg r1_xxreal_0 \\ X2 X0) \wedge (\neg r1_xxreal_0 X1 X2)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((\\ v1_xreal_0 X0) \Rightarrow ((k1_xxreal_3 (k3_xxreal_3 X1 X0) X0 = X1) \wedge (k3_xxreal_3 \\ (k1_xxreal_3 X1 X0) X0 = X1)))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (8)$$

Assume the following.

$$(m2_subset_1 np_0 k1_numbers k5_numbers) \wedge ((m1_subset_1 np_0 k5_numbers) \wedge (m1_subset_1 np_0 k1_numbers)) \quad (9)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (10)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge ((v1_xxreal_0 X1) \wedge (\neg v1_xreal_0 X1))) \Rightarrow (\neg v1_xreal_0 (k1_xxreal_3 X0 X1)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (v1_xxreal_0 (k3_xxreal_3 X0 X1)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (k1_xxreal_3 X0 X1 = k1_xxreal_3 X1 X0) \quad (14)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (15)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (16)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\neg (r1_xxreal_0 k6_numbers X0) \wedge ((\neg r1_xxreal_0 X1 X0) \wedge (\forall X2.(v1_xreal_0 X2) \Rightarrow (\neg (\neg r1_xxreal_0 X2 k6_numbers) \wedge (\neg r1_xxreal_0 X1 (k1_xxreal_3 X0 X2))))))))$$