

# t4\_binari\_3 (TMPquVtRpM- TUb5GVTtXf4e6a2USjxpA1NW1)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k5\_euclid : \iota \Rightarrow \iota$  be given. Let  $k3\_finseq\_2 : \iota \Rightarrow \iota$  be given. Let  $k6\_margrel1 : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_margrel1 : \iota$  be given. Let  $k1\_xboolean : \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_euclid : \iota \Rightarrow \iota$  be given. Let  $k13\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m2\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k4\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_euclid : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (2)$$

Assume the following.

$$k7\_margrel1 = k1\_xboolean \quad (3)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0)\wedge((v7\_ordinal1 X1)\wedge(m1\_subset\_1 X2 X0)))\Rightarrow(k5\_finseq\_2 X0 X1 X2 = k2\_finseq\_2 X1 X2) \quad (6)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow(k5\_euclid X0 = k4\_euclid X0) \quad (7)$$

Assume the following.

$$\forall X0.k3\_finseq\_2 X0 = k13\_finseq\_1 X0 \quad (8)$$

Assume the following.

$$\forall X0.\exists X1.(m1\_finseq\_1 X1 X0)\wedge((v1\_relat\_1 X1)\wedge((v4\_relat\_1 X1 k5\_numbers)\wedge((v5\_relat\_1 X1 X0)\wedge((v1\_funct\_1 X1)\wedge((v1\_xboole\_0 X1)\wedge((v1\_finset\_1 X1)\wedge(v1\_finseq\_1 X1))))))) \quad (9)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (10)$$

Assume the following.

$$\neg v1\_xboole\_0 k6\_margrel1 \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge((\neg v1\_xboole\_0 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))))\Rightarrow(\forall X2.(m2\_subset\_1 X2 X0 X1)\Rightarrow(m1\_subset\_1 X2 X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_2 X1 X0)\Rightarrow(\forall X2.(m2\_finseq\_2 X2 X0 X1)\Rightarrow(m2\_finseq\_1 X2 X0)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow((v1\_funct\_1 X1)\wedge((v1\_finseq\_1 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (14)$$

Assume the following.

$$m1\_subset\_1 k7\_margrel1 k6\_margrel1 \quad (15)$$

Assume the following.

$$m2\_subset\_1 k6\_numbers k1\_numbers k5\_numbers \quad (16)$$

Assume the following.

$$m1\_subset\_1 \ k5\_numbers \ (k1\_zfmisc\_1 \ k1\_numbers) \quad (17)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. ((\neg v1\_xboole\_0 \ X0) \wedge ((v7\_ordinal1 \\ X1) \wedge (m1\_subset\_1 \ X2 \ X0))) \Rightarrow (m2\_finseq\_2 \ (k5\_finseq\_2 \ X0 \ X1 \ X2) \\ X0 \ (k4\_finseq\_2 \ X1 \ X0)) \end{aligned} \quad (18)$$

Assume the following.

$$\forall X0. (v7\_ordinal1 \ X0) \Rightarrow (m2\_finseq\_2 \ (k5\_euclid \ X0) \ k1\_numbers \ (k1\_euclid \ X0)) \quad (19)$$

Assume the following.

$$\forall X0. \forall X1. (v7\_ordinal1 \ X0) \Rightarrow (m1\_finseq\_2 \ (k4\_finseq\_2 \ X0 \ X1) \ X1) \quad (20)$$

Assume the following.

$$\forall X0. (v7\_ordinal1 \ X0) \Rightarrow (m1\_finseq\_2 \ (k1\_euclid \ X0) \ k1\_numbers) \quad (21)$$

Assume the following.

$$\forall X0. (v7\_ordinal1 \ X0) \Rightarrow (k4\_euclid \ X0 = k5\_finseq\_2 \ k1\_numbers \ X0 \ k6\_numbers) \quad (22)$$

Assume the following.

$$k1\_xboolean = k6\_numbers \quad (23)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k13\_finseq\_1 \ X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (m2\_finseq\_1 \ X2 \ X0)) \quad (24)$$

Assume the following.

$$\forall X0. \forall X1. (v1\_xboole\_0 \ X0) \Rightarrow (\forall X2. (m1\_subset\_1 \ X2 \ (k1\_zfmisc\_1 \ (k2\_zfmisc\_1 \ X1 \ X0))) \Rightarrow (v1\_xboole\_0 \ X2)) \quad (25)$$

**Theorem 1**  $\forall X0. (v7\_ordinal1 \ X0) \Rightarrow (k5\_euclid \ X0 \in k3\_finseq\_2 \ k6\_margrel1).$