

t4_cqc_the3
(TMNkR2xCU4RUUUuF2WDX2ENUN4fTFNbHFM2)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r3_cqc_the1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_cqc_the1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow ((r1_tarski X1 X2) \Rightarrow (r1_tarski (k1_cqc_the1 \\ X0 X1) (k1_cqc_the1 X0 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\neg v1_xboole_0 (k3_cqc_lang X0)) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2 X0 X1) \Rightarrow (m1_subset_1 X2 X0)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (m1_subset_1 (k3_cqc_lang X0) (k1_zfmisc_1 \\ (k9_qc_lang1 X0))) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow (\forall X2.(m1_subset_1 X2 (k9_qc_lang1 \\ X0) \Rightarrow ((r3_cqc_the1 X0 X1 X2) \Leftrightarrow (X2 \in k1_cqc_the1 X0 X1)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1) \Leftrightarrow (\forall X2.(X2 \in X0) \Rightarrow \\ (X2 \in X1)) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (7)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow (\forall X3.(m1_subset_1 X3 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow (((r3_cqc_the1 X0 X2 X1) \wedge (r1_tarski X2 X3)) \Rightarrow \\ (r3_cqc_the1 X0 X3 X1)))))) \end{aligned}$$