

t4\_extreal1 (TM-  
SRN36CZrK5qdWVhE2FJn46ESMCcmQMWNQ)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k3\_extreal1 : \iota \Rightarrow \iota$  be given. Let  $k2\_supinf\_2 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k7\_numbers) \Rightarrow (((r1\_xxreal\_0 k6\_numbers X0) \Rightarrow (k3\_extreal1 X0 = X0)) \wedge ((\neg r1\_xxreal\_0 k6\_numbers X0) \Rightarrow (k3\_extreal1 X0 = k2\_supinf\_2 X0))) \quad (1)$$

**Theorem 1**

$$\forall X0.(m1\_subset\_1 X0 k7\_numbers) \Rightarrow ((\neg r1\_xxreal\_0 k6\_numbers X0) \Rightarrow (k3\_extreal1 X0 = k2\_supinf\_2 X0))$$