

t4\_fib\_num3 (TMN-  
QQDHLmwEnx39BXiPMoCZTu484dznDpa9)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_nat\_d : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k1\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7\_ordinal1 X2) \Rightarrow ((r1\_xxreal\_0 X0 X1) \Rightarrow (k7\_nat\_d (k2\_xcmplx\_0 \\ & X1 X2) X0 = k2\_nat\_1 (k7\_nat\_d X1 X0) X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.((\neg v1\_xboole\_0 X0) \wedge (v7\_ordinal1 X0)) \Rightarrow (\neg (X0 \neq np\_1) \wedge (r1\_xxreal\_0 X0 np\_1)) \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_2) \wedge (m2\_subset\_1 np\_2 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_2 k5\_numbers) \wedge (m1\_subset\_1 np\_2 k1\_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0) \wedge (m1\_subset\_1 X1 k5\_numbers)) \Rightarrow (k1\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0)\wedge(v1\_xxreal\_0 X1))\Rightarrow( (r1\_xxreal\_0 X0 X1)\vee(r1\_xxreal\_0 X1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow(v1\_xxreal\_0 X0) \quad (9)$$

**Theorem 1**

$$\forall X0.((\neg v1\_xboole\_0 X0)\wedge(v7\_ordinal1 X0))\Rightarrow(k2\_nat\_1 ( k7\_nat\_d X0 np\_1) np\_2 = k7\_nat\_d (k1\_nat\_1 X0 np\_2) np\_1)$$