

t4_funct_1
(TMGHafG7MDNstr35uvsRDN9YxVJ5DPYTDiJ)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.(X3 \in k9_xtuple_0 X0) \wedge (X2 = k1_funct_1 X0 X3)))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1.(X1 = k1_tarski X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (2)$$

Theorem 1

$$\forall X0. \forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((k9_xtuple_0 X1 = k1_tarski X0) \Rightarrow (k10_xtuple_0 X1 = k1_tarski (k1_funct_1 X1 X0)))$$