

t4_funct_7 (TMTisxvryt- BVvMMfR59WYGaGJKvYx96oPNk)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow \\ & (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow ((v2_funct_1 X2) \Rightarrow (\forall X3. \\ & (m1_subset_1 X3 X0) \Rightarrow (\forall X4.(m1_subset_1 X4 (k1_zfmisc_1 \\ & X0)) \Rightarrow ((k3_funct_2 X0 X1 X2 X3 \in k7_relset_1 X0 X1 X2 X4) \Rightarrow (X3 \in X4)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (k8_relset_1 X0 X1 X2 X3 = k8_relat_1 X2 X3) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k7_subset_1 X0 X1 X2 = k4_xboole_0 X1 X2) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0) \wedge \\ & (((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1)))))) \wedge (m1_subset_1 X3 X0))) \Rightarrow (k3_funct_2 X0 \\ & X1 X2 X3 = k1_funct_1 X2 X3) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(m1_subset_1 (k7_relset_1 X0 X1 X2 X3) (k1_zfmisc_1 X1)) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.\forall X2.(X2 = k8_relat_1 X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 \in k9_xtuple_0 X0)\wedge(k1_funct_1 X0 X3 \in X1)))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k4_xboole_0 X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 \in X0)\wedge(\neg X3 \in X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(\neg v1_xboole_0 X1)\Rightarrow \\ & (\forall X2.((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))))\Rightarrow((v2_funct_1 X2)\Rightarrow(\forall X3. \\ & (m1_subset_1 X3 X0)\Rightarrow(\forall X4.(m1_subset_1 X4 (k1_zfmisc_1 \\ & X0))\Rightarrow(\forall X5.(m1_subset_1 X5 (k1_zfmisc_1 X1))\Rightarrow((k3_funct_2 \\ & X0 X1 X2 X3 \in k7_subset_1 X1 (k7_relset_1 X0 X1 X2 X4) X5)\Rightarrow(X3 \in k7_subset_1 \\ & X0 X4 (k8_relset_1 X0 X1 X2 X5)))))))))) \end{aligned}$$