

t4_fuzzy_4

(TMRv9UcSe9M3exwULhRkPfNQH41vVBwBCq5)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_fuzzy_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_binop_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow \\ & (\forall X2.((v5_relat_1 X2 (k1_rcomp_1 k6_numbers np_1)) \wedge (\\ & (v1_funct_1 X2) \wedge ((v1_funct_2 X2 (k2_zfmisc_1 X0 X1) k1_numbers) \wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 (k2_zfmisc_1 X0 X1) \\ & k1_numbers)))))) \Rightarrow (\forall X3.(r1_xxreal_0 k6_numbers (k1_seq_1 \\ & X2 X3)) \wedge (r1_xxreal_0 (k1_seq_1 X2 X3) np_1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. \forall X4. ((\neg v1_xboole_0 \\ & X0) \wedge ((\neg v1_xboole_0 X1) \wedge ((v5_relat_1 X2 (k1_rcomp_1 k6_numbers \\ & np_1)) \wedge ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 (k2_zfmisc_1 X0 X1) \\ & k1_numbers) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1) k1_numbers)))))) \Rightarrow (k1_fuzzy_4 X0 X1 X2 X3 X4 = k1_binop_1 \\ & X2 X3 X4) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.k4_tarSKI X0 X1 = k2_tarSKI (k2_tarSKI X0 X1) (k1_tarSKI X0) \quad (4)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.\forall X2.k1_binop_1 X0 X1 X2 = k1_funct_1 X0 (k4_tarSKI X1 X2)) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v5_relat_1 X0 k1_numbers))\Rightarrow((v1_relat_1 X0)\wedge(v3_valued_0 X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(\neg v1_xboole_0 X1)\Rightarrow \\ & (\forall X2.((v5_relat_1 X2 (k1_rcomp_1 k6_numbers np_1))\wedge \\ & (v1_funct_1 X2)\wedge((v1_funct_2 X2 (k2_zfmisc_1 X0 X1) k1_numbers)\wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 (k2_zfmisc_1 X0 X1) \\ & k1_numbers))))))\Rightarrow(\forall X3.\forall X4.(r1_xxreal_0 k6_numbers \\ & (k1_fuzzy_4 X0 X1 X2 X3 X4))\wedge(r1_xxreal_0 (k1_fuzzy_4 X0 X1 X2 X3 \\ & X4) np_1)))) \end{aligned}$$