

t4_glib_000 (TM-
Gaax3HjJCP7d2WJZDXAhswohLZpSP84Qg)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_glib_000 : \iota \Rightarrow \iota$ be given. Let $k10_glib_000 : \iota \Rightarrow \iota$ be given. Let $k11_glib_000 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_glib_000 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (1)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0))))) \Rightarrow (\neg v1_xboole_0 (k6_glib_000 X0)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v5_relat_1 X1 X0)) \Rightarrow (m1_subset_1 (k2_relset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (4)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0))))) \Rightarrow ((v1_funct_1 (k11_glib_000 X0)) \wedge ((v1_funct_2 (k11_glib_000 X0) (k7_glib_000 X0) (k6_glib_000 X0)) \wedge (m1_subset_1 (k11_glib_000 X0) (k1_zfmisc_1 (k2_zfmisc_1 (k7_glib_000 X0) (k6_glib_000 X0))))) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow ((v1_funct_1 (k10_glib_000 \\ X0)) \wedge ((v1_funct_2 (k10_glib_000 X0) (k7_glib_000 X0) (k6_glib_000 \\ X0)) \wedge (m1_subset_1 (k10_glib_000 X0) (k1_zfmisc_1 (k2_zfmisc_1 \\ (k7_glib_000 X0) (k6_glib_000 X0)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow (((X1 \neq k1_xboole_0) \Rightarrow ((v1_funct_2 X2 X0 \\ X1) \Leftrightarrow (X0 = k1_relset_1 X0 X2))) \wedge ((X1 = k1_xboole_0) \Rightarrow ((v1_funct_2 \\ X2 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \end{aligned} \quad (9)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow ((k1_relset_1 (\\ k7_glib_000 X0) (k10_glib_000 X0) = k7_glib_000 X0) \wedge ((k1_relset_1 \\ (k7_glib_000 X0) (k11_glib_000 X0) = k7_glib_000 X0) \wedge ((r1_tarski \\ (k2_relset_1 (k6_glib_000 X0) (k10_glib_000 X0)) (k6_glib_000 \\ X0)) \wedge (r1_tarski (k2_relset_1 (k6_glib_000 X0) (k11_glib_000 \\ X0)) (k6_glib_000 X0)))))) \end{aligned}$$