

t4_goboard7 (TM-
cCZ9SVBQwSTVpJgmGHtk1UMxrxwuGzhuc)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $k3_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rltopsp1 : \iota \Rightarrow o$ be given. Let $l2_algstr_0 : \iota \Rightarrow o$ be given. Let $l2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_algstr_0 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_rltopsp1 : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 (k15_euclid X0))) \Rightarrow (\forall X2.(\\ m1_subset_1 X2 (u1_struct_0 (k15_euclid X0))) \Rightarrow (\forall X3.(m1_subset_1 & (1) \\ X3 (u1_struct_0 (k15_euclid X0))) \Rightarrow ((k3_rlvect_1 (k15_euclid \\ X0) X1 X2 = k3_rlvect_1 (k15_euclid X0) X3 X2) \Rightarrow (X1 = X3)))))) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 & (2) \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1\ X0) \Rightarrow & ((v2_pre_topc\ (k15_euclid\ X0)) \wedge \\ & ((v13_algstr_0\ (k15_euclid\ X0)) \wedge ((v2_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & ((v3_rlvect_1\ (k15_euclid\ X0)) \wedge ((v4_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & ((v5_rlvect_1\ (k15_euclid\ X0)) \wedge ((v6_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & ((v7_rlvect_1\ (k15_euclid\ X0)) \wedge ((v8_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & (v5_rltopsp1\ (k15_euclid\ X0)))))))))) \end{aligned} \quad (5)$$

Assume the following.

$$\neg v1_xboole_0\ k1_numbers \quad (6)$$

Assume the following.

$$\forall X0.(l2_algstr_0\ X0) \Rightarrow ((l2_struct_0\ X0) \wedge (l1_algstr_0\ X0)) \quad (7)$$

Assume the following.

$$\forall X0.(l1_rlvect_1\ X0) \Rightarrow (l2_algstr_0\ X0) \quad (8)$$

Assume the following.

$$\forall X0.(l1_rltopsp1\ X0) \Rightarrow ((l1_rlvect_1\ X0) \wedge (l1_pre_topc\ X0)) \quad (9)$$

Assume the following.

$$m1_subset_1\ k5_numbers\ (k1_zfmisc_1\ k1_numbers) \quad (10)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0) \Rightarrow ((v5_rltopsp1\ (k15_euclid\ X0)) \wedge (l1_rltopsp1\ (k15_euclid\ X0))) \quad (11)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2. & (((v2_rlvect_1\ X0) \wedge (l1_algstr_0 \\ & X0)) \wedge ((m1_subset_1\ X1\ (u1_struct_0\ X0)) \wedge (m1_subset_1\ X2\ (u1_struct_0 \\ & X0)))) \Rightarrow (k3_rlvect_1\ X0\ X1\ X2 = k3_rlvect_1\ X0\ X2\ X1) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1) \Rightarrow (v7_ordinal1\ X0) \quad (13)$$

Theorem 1

$$\begin{aligned} \forall X0.(m2_subset_1\ X0\ k1_numbers\ k5_numbers) \Rightarrow & (\forall X1. \\ & (m1_subset_1\ X1\ (u1_struct_0\ (k15_euclid\ X0))) \Rightarrow (\forall X2. \\ & m1_subset_1\ X2\ (u1_struct_0\ (k15_euclid\ X0))) \Rightarrow (\forall X3.(m1_subset_1 \\ & X3\ (u1_struct_0\ (k15_euclid\ X0))) \Rightarrow ((k3_rlvect_1\ (k15_euclid \\ & X0)\ X2\ X1 = k3_rlvect_1\ (k15_euclid\ X0)\ X2\ X3) \Rightarrow (X1 = X3)))) \end{aligned}$$