

t4_gobrd11 (TMUKssxNtrCfpB- MQdf4E3uvNN2iHLv6ZgWx)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k4_connsp_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. k3_xboole_0 X0 (k2_xboole_0 X1 X2) = k2_xboole_0 (k3_xboole_0 X0 X1) (k3_xboole_0 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_subset_1 X1 (k1_zfmisc_1 X0)) \wedge (m1_subset_1 X2 (k1_zfmisc_1 X0))) \Rightarrow (k4_subset_1 X0 X1 X2 = k2_xboole_0 X1 X2) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_subset_1 X1 (k1_zfmisc_1 X0)) \wedge (m1_subset_1 X2 (k1_zfmisc_1 X0))) \Rightarrow (m1_subset_1 (k4_subset_1 X0 X1 X2) (k1_zfmisc_1 X0)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((l1_pre_topc X0) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))))) \Rightarrow (m1_subset_1 (k4_connsp_3 X0 X1 X2) (k1_zfmisc_1 (u1_struct_0 (k1_pre_topc X0 X2)))) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.(l1_pre_topc\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1 \\ (u1_struct_0\ X0))) \Rightarrow (\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1 \\ (u1_struct_0\ X0))) \Rightarrow (k4_connsp_3\ X0\ X1\ X2 = k9_subset_1\ (u1_struct_0 \\ X0)\ X1\ X2))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0\ X0\ X1 = k3_xboole_0\ X1\ X0 \quad (7)$$

Theorem 1

$$\begin{aligned} \forall X0.((\neg v2_struct_0\ X0) \wedge ((v2_pre_topc\ X0) \wedge (l1_pre_topc \\ X0))) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ (u1_struct_0 \\ X0))) \Rightarrow (\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (u1_struct_0 \\ X0))) \Rightarrow (\forall X3.(m1_subset_1\ X3\ (k1_zfmisc_1\ (u1_struct_0 \\ X0))) \Rightarrow (k4_connsp_3\ X0\ (k4_subset_1\ (u1_struct_0\ X0)\ X1\ X2)\ X3 = \\ k4_subset_1\ (u1_struct_0\ (k1_pre_topc\ X0\ X3))\ (k4_connsp_3\ X0 \\ X1\ X3)\ (k4_connsp_3\ X0\ X2\ X3)))))) \end{aligned}$$