

# t4\_nat\_2 (TMNMNUBbLZ- MAvqG863pWCKy9mUv3uRbmXqu)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k3\_nat.d : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k3\_xcmplx\_0 np\_1 X0 = X0) \quad (2)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((X0 \neq k6\_numbers) \Rightarrow (k3\_nat.d (k3\_xcmplx\_0 X0 X1) X0 = X1))) \quad (3)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (4)$$

Assume the following.

$$v1\_xboole\_0 np\_0 \quad (5)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (6)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(v7\_ordinal1\ X0)\Rightarrow(v1\_xcmplx\_0\ X0) \quad (9)$$

**Theorem 1**  $\forall X0.(v7\_ordinal1\ X0)\Rightarrow(k3\_nat\_d\ X0\ np\_1 = X0).$