

t4_prvect_3 (TMK-
tjdPMW9ikoNSCtqN6n5CPKZUZLfgZQew)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_card_3 : \iota \Rightarrow \iota$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $v2_funct_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $np_2 : \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \neg (X0 \in X1) \wedge (v1_xboole_0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((X1 = k9_finseq_1 X0) \Leftrightarrow ((k3_finseq_1 X1 = np_1) \wedge (k1_funct_1 X1 np_1 = X0))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((X1 = k9_finseq_1 X0) \Leftrightarrow ((k4_finseq_1 X1 = k2_finseq_1 np_1) \wedge (k10_xtuple_0 X1 = k1_tarski X0))) \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (\\ v1_funct_1 X1)) \Rightarrow (\neg(k9_xtuple_0 X1 = k6_domain_1 k5_numbers np_1) \wedge \\ ((k1_funct_1 X1 np_1 = X0) \wedge (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\ X2 X0 (k4_card_3 X1)) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ X0 (k4_card_3 X1)))))) \Rightarrow (\neg(v2_funct_1 X2) \wedge ((v2_funct_2 X2 (k4_card_3 \\ X1)) \wedge (\forall X3.(X3 \in X0) \Rightarrow (k1_funct_1 X2 X3 = k9_finseq_1 X3)))))))))) \end{aligned} \quad (4)$$

Assume the following.

$$(k2_finseq_1 np_1 = k1_tarski np_1) \wedge (k2_finseq_1 np_2 = k2_tarski np_1 np_2) \quad (5)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.k9_finseq_1 X0 = k5_finseq_1 X0 \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow \\ (k6_domain_1 X0 X1 = k1_tarski X1) \end{aligned} \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (k4_finseq_1 X0 = k9_xtuple_0 X0) \end{aligned} \quad (10)$$

Assume the following.

$$\forall X0.v1_finseq_1 (k5_finseq_1 X0) \quad (11)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (12)$$

Assume the following.

$$\forall X0.(v1_relat_1 (k5_finseq_1 X0)) \wedge (v1_funct_1 (k5_finseq_1 X0)) \quad (13)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\exists X1.((v1_funct_1 X1) \wedge (\\ (v1_funct_2 X1 X0 (k4_card_3 (k9_finseq_1 X0))) \wedge (m1_subset_1 \\ X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 (k4_card_3 (k9_finseq_1 X0)))))) \wedge \\ ((v2_funct_1 X1) \wedge ((v2_funct_2 X1 (k4_card_3 (k9_finseq_1 X0))) \wedge \\ (\forall X2.(X2 \in X0) \Rightarrow (k1_funct_1 X1 X2 = k9_finseq_1 X2)))))) \end{aligned}$$