

t4_recdef_2
(TMZD4jPRqp8Y2495HwQsjebNghC2qZgtZEs)

October 27, 2020

Let $k6_xtuple_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k8_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v3_xtuple_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. (v3_xtuple_0 X0) \Rightarrow (k6_xtuple_0 (k7_xtuple_0 X0) (k8_xtuple_0 X0) (k5_xtuple_0 X0) (k2_xtuple_0 X0) = X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. v3_xtuple_0 (k6_xtuple_0 X0 X1 X2 X3) \quad (2)$$

Theorem 1

$$\forall X0. (\exists X1. \exists X2. \exists X3. \exists X4. X0 = k6_xtuple_0 X1 X2 X3 X4) \Rightarrow (X0 = k6_xtuple_0 (k7_xtuple_0 X0) (k8_xtuple_0 X0) (k5_xtuple_0 X0) (k2_xtuple_0 X0))$$