

t4_rfinseq2
(TMZK3PK9ANfRF1sbTn6d8QK2DDFNzWGuzT)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_rfinseq2 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k4_rfinseq2 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v2_membered : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((X1 = k9_finseq_1 X0) \Leftrightarrow ((k3_finseq_1 X1 = np_1) \wedge (k1_funct_1 X1 np_1 = X0))) \quad (1)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))))) \quad (2)$$

Assume the following.

$$\forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1)) \quad (3)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Leftrightarrow(m1_finseq_1 X1 X0) \quad (5)$$

Assume the following.

$$\forall X0.k9_finseq_1 X0 = k5_finseq_1 X0 \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow (k12_finseq_1 X0 X1 = k5_finseq_1 X1) \quad (11)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (12)$$

Assume the following.

$$\forall X0.(v1_relat_1 (k5_finseq_1 X0)\wedge(v1_funct_1 (k5_finseq_1 X0))) \quad (13)$$

Assume the following.

$$v3_membered k1_numbers \quad (14)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (16)$$

Assume the following.

$$\forall X0.(m1_finseq_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k2_rfinseq2 X0) k5_numbers) \quad (17)$$

Assume the following.

$$\forall X0.(m2_finseq_1 X0 k1_numbers) \Rightarrow (k4_rfinseq2 X0 = k1_seq_1 X0 (k2_rfinseq2 X0)) \quad (18)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k5_numbers) \Rightarrow ((X1 = k2_rfinseq2 X0) \Leftrightarrow (((k3_finseq_1 X0 = k6_numbers) \Rightarrow \\ & (X1 = k6_numbers)) \wedge ((\neg r1_xxreal_0 (k3_finseq_1 X0) k6_numbers) \Rightarrow \\ & ((X1 \in k4_finseq_1 X0) \wedge (\forall X2.(m1_subset_1 X2 k5_numbers) \Rightarrow \\ & (\forall X3.(m1_subset_1 X3 k1_numbers) \Rightarrow (\forall X4.(m1_subset_1 X4 k1_numbers) \Rightarrow (((X2 \in k4_finseq_1 X0) \wedge ((X3 = k1_seq_1 X0 X2) \wedge \\ & (X4 = k1_seq_1 X0 X1))) \Rightarrow (r1_xxreal_0 X4 X3)))))) \wedge (\forall X2.(m1_subset_1 X2 k5_numbers) \Rightarrow (((X2 \in k4_finseq_1 X0) \wedge (k1_seq_1 X0 X2 = k1_seq_1 X0 X1)) \Rightarrow (r1_xxreal_0 X1 X2)))))))))) \end{aligned} \quad (19)$$

Assume the following.

$$\forall X0.(v3_membered X0) \Rightarrow (v2_membered X0) \quad (20)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xxreal_0 X0) \quad (21)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (22)$$

Assume the following.

$$\forall X0.\forall X1.(v3_membered X1) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v3_valued_0 X2)) \quad (23)$$

Assume the following.

$$\forall X0.(v6_membered X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow (v7_ordinal1 X1)) \quad (24)$$

Assume the following.

$$\forall X0.(v2_membered X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow (v1_xxreal_0 X1)) \quad (25)$$

Theorem 1

$$\forall X0.(m2_finseq_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow ((X0 = k12_finseq_1 k1_numbers X1) \Rightarrow ((k2_rfinseq2 X0 = np_1) \wedge (k4_rfinseq2 X0 = X1))))$$