

t4_rlvect_1
(TMP9ZmfZmPGa9PE2H3NN6zAZMpAuN1H57GK)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $l2_algstr_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_struct_0 : \iota \Rightarrow \iota$ be given. Let $l2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_algstr_0 : \iota \Rightarrow o$ be given. Let $v10_algstr_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. ((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v3_rlvect_1 \\ X0) \wedge ((v4_rlvect_1 X0) \wedge (l2_algstr_0 X0)))))) \Rightarrow (\forall X1. (m1_subset_1 \\ X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_subset_1 X2 (u1_struct_0 \\ X0)) \Rightarrow ((k1_algstr_0 X0 X1 X2 = k4_struct_0 X0) \Rightarrow (k1_algstr_0 X0 X2 \\ X1 = k4_struct_0 X0)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. (l2_algstr_0 X0) \Rightarrow ((l2_struct_0 X0) \wedge (l1_algstr_0 X0)) \quad (2)$$

Assume the following.

$$\forall X0. (l2_algstr_0 X0) \Rightarrow ((v4_rlvect_1 X0) \Leftrightarrow (\forall X1. (\\ m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (k1_algstr_0 X0 X1 (k4_struct_0 \\ X0) = X1)))) \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0. (l1_algstr_0 X0) \Rightarrow ((v3_rlvect_1 X0) \Leftrightarrow (\forall X1. (\\ m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_subset_1 X2 \\ (u1_struct_0 X0)) \Rightarrow (\forall X3. (m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow \\ (k1_algstr_0 X0 (k1_algstr_0 X0 X1 X2) X3 = k1_algstr_0 X0 X1 (k1_algstr_0 \\ X0 X2 X3)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0. (l2_algstr_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 \\ X0)) \Rightarrow ((v10_algstr_0 X1 X0) \Leftrightarrow (\exists X2. (m1_subset_1 X2 (u1_struct_0 \\ X0)) \wedge (k1_algstr_0 X0 X1 X2 = k4_struct_0 X0)))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v13_algstr_0 X0) \wedge (l2_algstr_0 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (v10_algstr_0 X1 X0)) \end{aligned} \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v3_rlvect_1 \\ & X0) \wedge ((v4_rlvect_1 X0) \wedge (l2_algstr_0 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 (u1_struct_0 X0)) \Rightarrow ((k1_algstr_0 X0 X1 (k4_struct_0 X0) = X1) \wedge \\ & (k1_algstr_0 X0 (k4_struct_0 X0) X1 = X1))) \end{aligned}$$