

# t4\_rlvect\_2 (TMGGCcqNKKhk- LEmd9oniY1GePgYowF9uWmw)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l2\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_algstr\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_rlvect\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_finseq\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X2))) \Rightarrow (m1\_subset\_1 X0 X2) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge (l2\_algstr\_0 X0)))))) \Rightarrow \\ & (\forall X1. (m2\_finseq\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (\forall X2. (m2\_finseq\_1 X2 (u1\_struct\_0 X0)) \Rightarrow ((k3\_finseq\_1 X1 = k3\_finseq\_1 X2) \wedge (\forall X3. \\ & (m2\_subset\_1 X3 k1\_numbers k5\_numbers) \Rightarrow (\forall X4. (m1\_subset\_1 X4 (u1\_struct\_0 X0)) \Rightarrow ((X3 \in k4\_finseq\_1 X1) \wedge (X4 = k1\_funct\_1 X2 X3)) \Rightarrow (k1\_funct\_1 X1 X3 = k4\_algstr\_0 X0 X4)))))) \Rightarrow (k4\_rlvect\_1 X0 X1 = k4\_algstr\_0 X0 (k4\_rlvect\_1 X0 X2)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow((v1\_funct\_1 X1)\wedge((v1\_finseq\_1 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (5)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow(m1\_subset\_1 (k4\_finseq\_1 X0) (k1\_zfmisc\_1 k5\_numbers)) \quad (6)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow(m2\_subset\_1 (k3\_finseq\_1 X0) k1\_numbers k5\_numbers) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge((v5\_relat\_1 X1 X0)\wedge(v1\_funct\_1 X1)))\Rightarrow(\forall X2.(X2 \in k9\_xtuple\_0 X1)\Rightarrow(k7\_partfun1 X0 X1 X2 = k1\_funct\_1 X1 X2)) \quad (8)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow(\forall X1.(m2\_subset\_1 X1 k1\_numbers k5\_numbers)\Rightarrow((X1 = k3\_finseq\_1 X0)\Leftrightarrow(k2\_finseq\_1 X1 = k9\_xtuple\_0 X0))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (11)$$

### Theorem 1

$$\forall X0.((\neg v2\_struct\_0 X0)\wedge((v13\_algstr\_0 X0)\wedge((v2\_rlvect\_1 X0)\wedge((v3\_rlvect\_1 X0)\wedge((v4\_rlvect\_1 X0)\wedge(l2\_algstr\_0 X0))))))\Rightarrow(\forall X1.(m2\_finseq\_1 X1 (u1\_struct\_0 X0))\Rightarrow(\forall X2.(m2\_finseq\_1 X2 (u1\_struct\_0 X0))\Rightarrow(((k3\_finseq\_1 X1 = k3\_finseq\_1 X2)\wedge(\forall X3.(m1\_subset\_1 X3 k5\_numbers)\Rightarrow((X3 \in k1\_relset\_1 k5\_numbers X1)\Rightarrow(k1\_funct\_1 X2 X3 = k4\_algstr\_0 X0 (k7\_partfun1 (u1\_struct\_0 X0) X1 X3))))))\Rightarrow(k4\_rlvect\_1 X0 X2 = k4\_algstr\_0 X0 (k4\_rlvect\_1 X0 X1))))))$$