

t4_scmpds_4
(TMMcjPikvJi5m7C2Mg3QzfeJrvn63US12uk)

October 27, 2020

Let $v1_ami_2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_scmpds_2 : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k2_scmpds_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k2_ami_2 : \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_int_1 : \iota \Rightarrow o$ be given. Let $k1_int_2 : \iota \Rightarrow \iota$ be given. Let $k16_complex1 : \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 X1) \wedge ((\neg v3_xxreal_0 X1) \wedge (\neg v2_xxreal_0 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0.\neg(X0 \in k2_ami_2) \wedge (\forall X1.(m1_subset_1 X1 k5_numbers) \Rightarrow (X0 \neq k4_tarski np_1 X1)) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k2_xcmplx_0 X0 k6_numbers = X0) \quad (4)$$

Assume the following.

$$(m2_subset_1 np_0 k1_numbers k5_numbers) \wedge ((m1_subset_1 np_0 k5_numbers) \wedge (m1_subset_1 np_0 k1_numbers)) \quad (5)$$

Assume the following.

$$v1_xboole_0 \text{ np_}0 \quad (6)$$

Assume the following.

$$k4_xcmplx_0 \text{ np_}0 = \text{np_}0 \quad (7)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0.(v1_int_1 X0) \Rightarrow (k1_int_2 X0 = k16_complex1 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k4_xcmplx_0 (k4_xcmplx_0 X0) = X0) \quad (11)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((v1_xcmplx_0 (k4_xcmplx_0 X0)) \wedge (v1_xreal_0 (k4_xcmplx_0 X0))) \quad (12)$$

Assume the following.

$$\forall X0.(v1_int_1 X0) \Rightarrow ((v1_xcmplx_0 (k4_xcmplx_0 X0)) \wedge (v1_int_1 (k4_xcmplx_0 X0))) \quad (13)$$

Assume the following.

$$\forall X0.((\neg v3_xreal_0 X0) \wedge (v1_xreal_0 X0)) \Rightarrow ((v1_xcmplx_0 (k4_xcmplx_0 X0)) \wedge (\neg v2_xreal_0 (k4_xcmplx_0 X0))) \quad (14)$$

Assume the following.

$$\forall X0.(v1_int_1 X0) \Rightarrow (m1_subset_1 (k1_int_2 X0) k5_numbers) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.k4_tarski X0 X1 = k2_tarski (k2_tarski X0 X1) (k1_tarski X0) \quad (16)$$

Assume the following.

$$\forall X0.(v1_int_1 X0) \Rightarrow (\forall X1.(v1_int_1 X1) \Rightarrow (k2_scmpds_2 X0 X1 = k4_tarski \text{ np_}1 (k1_int_2 (k2_xcmplx_0 X0 X1)))) \quad (17)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (((r1_xxreal_0 k6_numbers X0) \Rightarrow (k16_complex1 X0 = X0)) \wedge ((\neg r1_xxreal_0 k6_numbers X0) \Rightarrow (k16_complex1 X0 = k4_xcplx_0 X0))) \quad (18)$$

Assume the following.

$$\forall X0.(v1_ami_2 X0) \Leftrightarrow (X0 \in k2_ami_2) \quad (19)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (20)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcplx_0 X0) \quad (21)$$

Assume the following.

$$\forall X0.(v1_int_1 X0) \Rightarrow (v1_xreal_0 X0) \quad (22)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\neg v3_xxreal_0 X0) \quad (23)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_int_1 X0) \quad (24)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (25)$$

Theorem 1

$$\forall X0.((v1_ami_2 X0) \wedge (m1_subset_1 X0 (u1_struct_0 k1_scmpds_2))) \Rightarrow (\exists X1.(m1_subset_1 X1 k5_numbers) \wedge (X0 = k2_scmpds_2 X1 k6_numbers))$$