

t4_semi_af1

(TMYTqhL65AEbzS6Rf6qUhqPMC76gSh9mZa9)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v1_semi_af1 : \iota \Rightarrow o$ be given. Let $l1_analoaf : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r2_analoaf : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the follow-

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_semi_af1 X0) \wedge (l1_analoaf \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 \\ & (u1_struct_0 X0)) \Rightarrow (\forall X4.(m1_subset_1 X4 (u1_struct_0 X0)) \Rightarrow \\ & ((r2_analoaf X0 X1 X2 X3 X4) \Rightarrow (r2_analoaf X0 X2 X1 X3 X4)))))) \end{aligned}$$