

t4_setlim_1 (TMKosqFECqR- rDHu2pnbRwqkW9TneKc8VmtR)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X3) \wedge \\ & ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))))) \Rightarrow (\neg(X2 \in k2_relset_1 X1 X3) \wedge (\forall X4. \neg(X4 \in X0) \wedge (k1_funct_1 \\ & X3 X4 = X2))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (\neg v1_xboole_0 X1) \Rightarrow \\ & (\forall X2. (m1_subset_1 X2 X0) \Rightarrow (\forall X3. ((v1_funct_1 X3) \wedge \\ & ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))))) \Rightarrow (k3_funct_2 X0 X1 X3 X2 \in k2_relset_1 X1 X3)))) \end{aligned} \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((\neg v1_xboole_0 X0) \wedge \\ & (((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1)))))) \wedge (m1_subset_1 X3 X0))) \Rightarrow (k3_funct_2 X0 \\ & X1 X2 X3 = k1_funct_1 X2 X3) \end{aligned} \quad (5)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (\neg v1_xboole_0 \ X1) \Rightarrow (\forall X2. ((v1_funct_1 \\ & X2) \wedge ((v1_funct_2 \ X2 \ k5_numbers \ X1) \wedge (m1_subset_1 \ X2 \ (k1_zfmisc_1 \\ & (k2_zfmisc_1 \ k5_numbers \ X1)))))) \Rightarrow ((X0 \in k2_relset_1 \ X1 \ X2) \Leftrightarrow (\exists X3. \\ & (m1_subset_1 \ X3 \ k5_numbers) \wedge (X0 = k3_funct_2 \ k5_numbers \ X1 \ X2 \ X3)))) \end{aligned}$$