

t4_subset_1
(TMKFbquBjS1tvpfDEvYtZJf9wR6x4Jgnz3Y)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \neg(X0 \neq k1_xboole_0) \wedge (\forall X1. \neg X1 \in X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. (X2 \in X1) \Rightarrow (X2 \in X0)) \quad (2)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Leftrightarrow (\forall X1. \neg X1 \in X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \Rightarrow ((m1_subset_1 X1 X0) \Leftrightarrow (X1 \in X0))) \wedge ((v1_xboole_0 X0) \Rightarrow ((m1_subset_1 X1 X0) \Leftrightarrow (v1_xboole_0 X1))) \quad (4)$$

Theorem 1

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\neg(X1 \neq k1_xboole_0) \wedge (\forall X2. (m1_subset_1 X2 X0) \Rightarrow (\neg X2 \in X1)))$$