

t4_topreal3

(TMN3wmBxKkFPPFftuGFC1ZB5R6proo97VMc)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k1_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 \\ (k15_euclid np_2))) \Rightarrow (k1_rlvect_1 (k15_euclid np_2) X1 X0 = k19_euclid \\ (k4_real_1 X0 (k17_euclid X1)) (k4_real_1 X0 (k18_euclid X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid \\ (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((v1_xreal_0 X0) \wedge (m1_subset_1 X1 k1_numbers)) \Rightarrow \\ (m1_subset_1 (k4_real_1 X0 X1) k1_numbers) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ (m1_subset_1 (k18_euclid X0) k1_numbers) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ (m1_subset_1 (k17_euclid X0) k1_numbers) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (6)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid (k1_rlvect_1 (k15_euclid \\ np_2) X0 X1) = k4_real_1 X1 (k17_euclid X0)) \wedge (k18_euclid (k1_rlvect_1 \\ (k15_euclid np_2) X0 X1) = k4_real_1 X1 (k18_euclid X0)))) \end{aligned}$$