

t4_waybel_0
(TMJ7rTorEyDbv7hQp7xSycMwPaBTABUB6RM)

October 27, 2020

Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $g1_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $u1_orders_2 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v2_waybel_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r2_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. (l1_orders_2 X0) \Rightarrow (\forall X1. (l1_orders_2 X1) \Rightarrow ((\\ g1_orders_2 (u1_struct_0 X0) (u1_orders_2 X0) = g1_orders_2 (u1_struct_0 \\ X1) (u1_orders_2 X1)) \Rightarrow (\forall X2. (m1_subset_1 X2 (u1_struct_0 \\ X0)) \Rightarrow (\forall X3. (m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow (\forall X4. \\ (m1_subset_1 X4 (u1_struct_0 X1)) \Rightarrow (\forall X5. (m1_subset_1 X5 \\ (u1_struct_0 X1)) \Rightarrow (((X2 = X4) \wedge (X3 = X5)) \Rightarrow (((r1_orders_2 X0 X2 X3) \Rightarrow \\ (r1_orders_2 X1 X4 X5)) \wedge ((r2_orders_2 X0 X2 X3) \Rightarrow (r2_orders_2 X1 \\ X4 X5)))))))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. (l1_orders_2 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\ (u1_struct_0 X0))) \Rightarrow ((v2_waybel_0 X1 X0) \Leftrightarrow (\forall X2. (m1_subset_1 \\ X2 (u1_struct_0 X0)) \Rightarrow (\forall X3. (m1_subset_1 X3 (u1_struct_0 \\ X0)) \Rightarrow (\neg(X2 \in X1) \wedge ((X3 \in X1) \wedge (\forall X4. (m1_subset_1 X4 (u1_struct_0 \\ X0)) \Rightarrow (\neg(X4 \in X1) \wedge ((r1_orders_2 X0 X4 X2) \wedge (r1_orders_2 X0 X4 X3)))))))))) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} \forall X0. (l1_orders_2 X0) \Rightarrow (\forall X1. (l1_orders_2 X1) \Rightarrow ((\\ g1_orders_2 (u1_struct_0 X0) (u1_orders_2 X0) = g1_orders_2 (u1_struct_0 \\ X1) (u1_orders_2 X1)) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ (u1_struct_0 X0))) \Rightarrow (\forall X3. (m1_subset_1 X3 (k1_zfmisc_1 \\ (u1_struct_0 X1))) \Rightarrow (((X2 = X3) \wedge (v2_waybel_0 X2 X0)) \Rightarrow (v2_waybel_0 \\ X3 X1)))))) \end{aligned}$$